


Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application.

Listing of Claims

1. (Currently Amended) A method for treating pain in a patient, comprising administering to a patient ~~in need thereof~~ experiencing pain an effective amount of a JNK Inhibitor or a pharmaceutically acceptable salt, solvate or stereoisomer thereof.
2. (Currently Amended) A method for treating pain in a patient, comprising administering to a patient ~~in need thereof~~ experiencing pain an effective amount of a compound having the following formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

wherein:

A is a direct bond, $-(CH_2)_a-$, $-(CH_2)_bCH=CH(CH_2)_c-$, or $-(CH_2)_bC\equiv C(CH_2)_c-$;

R₁ is aryl, heteroaryl or heterocycle fused to phenyl, each being optionally substituted with one to four substituents independently from R₃;

R₂ is -R₃, -R₄, $-(CH_2)_bC(=O)R_5$, $-(CH_2)_bC(=O)OR_5$, $-(CH_2)_bC(=O)NR_5R_6$, $-(CH_2)_bC(=O)NR_5(CH_2)_cC(=O)R_6$, $-(CH_2)_bNR_5C(=O)R_6$, $-(CH_2)_bNR_5C(=O)NR_6R_7$, $-(CH_2)_bNR_5R_6$, $-(CH_2)_bOR_5$, $-(CH_2)_bSO_dR_5$ or $-(CH_2)_bSO_2NR_5R_6$;

a is 1, 2, 3, 4, 5 or 6;

b and c are the same or different and at each occurrence independently 0, 1, 2, 3 or 4;

d is at each occurrence 0, 1 or 2;


R₃ is at each occurrence independently halogen, hydroxy, carboxy, alkyl, alkoxy, haloalkyl, acyloxy, thioalkyl, sulfinylalkyl, sulfonylalkyl, hydroxyalkyl, aryl, substituted aryl, arylalkyl,

heterocycle, heterocycloalkyl, $-C(=O)OR_8$, $-OC(=O)R_8$, $-C(=O)NR_8R_9$, $-C(=O)NR_8OR_9$, $-SO_2NR_8R_9$, $-NR_8SO_2R_9$, $-CN$, $-NO_2$, $-NR_8R_9$, $-NR_8C(=O)R_9$, $-NR_8C(=O)(CH_2)_bOR_9$, $-NR_8C(=O)(CH_2)_bR_9$, $-O(CH_2)_bNR_8R_9$, or heterocycle fused to phenyl;

R_4 is alkyl, aryl, arylalkyl, heterocycle or heterocycloalkyl, each being optionally substituted with one to four substituents independently from R_3 , or R_4 is halogen or hydroxy;

R_5 , R_6 and R_7 are the same or different and at each occurrence independently hydrogen, alkyl, aryl, arylalkyl, heterocycle or heterocycloalkyl, wherein each of R_5 , R_6 and R_7 are optionally substituted with one to four substituents independently selected from R_3 ; and R_8 and R_9 are the same or different and at each occurrence independently hydrogen, alkyl, aryl, arylalkyl, heterocycle, or heterocycloalkyl, or R_8 and R_9 taken together with the atom or atoms to which they are bonded form a heterocycle, wherein each of R_8 , R_9 , and R_8 and R_9 taken together to form a heterocycle are optionally substituted with one to four substituents independently selected from R_3 .

3. (Withdrawn) A method for treating, preventing, managing and/or modifying pain in a patient, comprising administering to a patient in need thereof an effective amount of a compound having the following formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

wherein:

R_1 is aryl or heteroaryl optionally substituted with one to four substituents independently selected from R_7 ;

R_2 is hydrogen;

R_3 is hydrogen or lower alkyl;

R₄ represents one to four optional substituents, wherein each substituent is the same or different and independently halogen, hydroxy, lower alkyl or lower alkoxy;

R₅ and R₆ are the same or different and independently -R₈, -(CH₂)_aC(=O)R₉, -(CH₂)_aC(=O)OR₉, -(CH₂)_aC(=O)NR₉R₁₀, -(CH₂)_aC(=O)NR₉(CH₂)_bC(=O)R₁₀, -(CH₂)_aNR₉C(=O)R₁₀, (CH₂)_aNR₁₁C(=O)NR₉R₁₀, -(CH₂)_aNR₉R₁₀, -(CH₂)_aOR₉, -(CH₂)_aSO_cR₉ or -(CH₂)_aSO₂NR₉R₁₀;

or R₅ and R₆ taken together with the nitrogen atom to which they are attached to form a heterocycle or substituted heterocycle;

R₇ is at each occurrence independently halogen, hydroxy, cyano, nitro, carboxy, alkyl, alkoxy, haloalkyl, acyloxy, thioalkyl, sulfinylalkyl, sulfonylalkyl, hydroxyalkyl, aryl, arylalkyl, heterocycle, heterocycloalkyl, -C(=O)OR₈, -OC(=O)R₈, -C(=O)NR₈R₉, -C(=O)NR₈OR₉, -SO_cR₈, -SO_cNR₈R₉, -NR₈SO_cR₉, -NR₈R₉, -NR₈C(=O)R₉, -NR₈C(=O)(CH₂)_bOR₉, -NR₈C(=O)(CH₂)_bR₉, -O(CH₂)_bNR₈R₉, or heterocycle fused to phenyl;


R₈, R₉, R₁₀ and R₁₁ are the same or different and at each occurrence independently hydrogen, alkyl, substituted alkyl, aryl, arylalkyl, heterocycle or heterocycloalkyl.;

or R₈ and R₉ taken together with the atom or atoms to which they are attached to form a heterocycle;

a and *b* are the same or different and at each occurrence independently 0, 1, 2, 3 or 4; and

c is at each occurrence 0, 1 or 2.

4. (Withdrawn) A method for treating, preventing, managing and/or modifying pain in a patient, comprising administering to a patient in need thereof an effective amount of a compound having the following formula:




or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

wherein R_0 is -O-, -S-, -S(O)-, -S(O)₂-, NH or -CH₂-;

the compound being (i) unsubstituted, (ii) monosubstituted and having a first substituent, or (iii) disubstituted and having a first substituent and a second substituent;

the first or second substituent, when present, is at the 3, 4, 5, 7, 8, 9, or 10 position, wherein the first and second substituent, when present, are independently alkyl, hydroxy, halogen, nitro, trifluoromethyl, sulfonyl, carboxyl, alkoxycarbonyl, alkoxy, aryl, aryloxy, arylalkyloxy, arylalkyl, cycloalkylalkyloxy, cycloalkyloxy, alkoxyalkyl, alkoxyalkoxy, aminoalkoxy, mono-alkylaminoalkoxy, di-alkylaminoalkoxy, or a group represented by




formula (a), (b), (c), (d), (e), or (f):

wherein R_3 and R_4 are taken together and represent alkylidene or a heteroatom-containing cyclic alkylidene or R_3 and R_4 are independently hydrogen, alkyl, cycloalkyl, aryl, arylalkyl, cycloalkylalkyl, aryloxyalkyl, alkoxyalkyl, aminoalkyl, mono-alkylaminoalkyl, or di-alkylaminoalkyl; and

R₅ is hydrogen, alkyl, cycloalkyl, aryl, arylalkyl, cycloalkylalkyl, alkoxy, alkoxyalkyl, alkoxycarbonylalkyl, amino, mono-alkylamino, di-alkylamino, arylamino, arylalkylamino, cycloalkylamino, cycloalkylalkylamino, aminoalkyl, mono-alkylaminoalkyl, or di-alkylaminoalkyl.

5. (Withdrawn) The method of claim 2 wherein A is a direct bond.
6. (Withdrawn) The method of claim 2 wherein A is $-(CH_2)_a-$.
7. (Withdrawn) The method of claim 2 wherein A is $-(CH_2)_bCH=CH(CH_2)_c-$.
8. (Withdrawn) The method of claim 2 wherein A is $-(CH_2)_bC\equiv C(CH_2)_c-$.
9. (Withdrawn) The method of claim 2 wherein the compound has the following formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

wherein:

A is a direct bond, $-(CH_2)_a-$, $-(CH_2)_bCH=CH(CH_2)_c-$, or $-(CH_2)_bC\equiv C(CH_2)_c-$;

R₁ is aryl, heteroaryl or heterocycle fused to phenyl, each being optionally substituted with one to four substituents independently from R₃;

R₂ is -R₃, -R₄, $-(CH_2)_bC(=O)R_5$, $-(CH_2)_bC(=O)OR_5$,

$-(CH_2)_bC(=O)NR_5R_6$, $-(CH_2)_bC(=O)NR_5(CH_2)_cC(=O)R_6$, $-(CH_2)_bNR_5C(=O)R_6$,

$-(CH_2)_bNR_5C(=O)NR_6R_7$, $-(CH_2)_bNR_5R_6$, $-(CH_2)_bOR_5$, $-(CH_2)_bSO_2R_5$ or $-(CH_2)_bSO_2NR_5R_6$;

a is 1, 2, 3, 4, 5 or 6;

b and c are the same or different and at each occurrence independently selected from 0, 1, 2, 3 or 4;

d is at each occurrence 0, 1 or 2;


R_3 is at each occurrence independently halogen, hydroxy, carboxy, alkyl, alkoxy, haloalkyl, acyloxy, thioalkyl, sulfinylalkyl, sulfonylalkyl, hydroxyalkyl, aryl, arylalkyl, heterocycle, heterocycloalkyl, $-C(=O)OR_8$, $-OC(=O)R_8$, $-C(=O)NR_8R_9$, $-C(=O)NR_8OR_9$, $-SO_2NR_8R_9$, $-NR_8SO_2R_9$, $-CN$, $-NO_2$, $-NR_8R_9$, $-NR_8C(=O)R_9$, $-NR_8C(=O)(CH_2)_bOR_9$, $-NR_8C(=O)(CH_2)_bR_9$, $-O(CH_2)_bNR_8R_9$, or heterocycle fused to phenyl;

R_4 is alkyl, aryl, arylalkyl, heterocycle or heterocycloalkyl, each being optionally substituted with one to four substituents independently from R_3 , or R_4 is halogen or hydroxy;

R_5 , R_6 and R_7 are the same or different and at each occurrence independently hydrogen, alkyl, aryl, arylalkyl, heterocycle or heterocycloalkyl, wherein each of R_5 , R_6 and R_7 are optionally substituted with one to four substituents independently selected from R_3 ; and

R_8 and R_9 are the same or different and at each occurrence independently hydrogen, alkyl, aryl, arylalkyl, heterocycle, or heterocycloalkyl, or R_8 and R_9 taken together with the atom or atoms to which they are bonded form a heterocycle, wherein each of R_8 , R_9 , and R_8 and R_9 taken together to form a heterocycle are optionally substituted with one to four substituents independently selected from R_3 .

10. (Withdrawn) The method of claim 2 wherein the compound has the following formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

wherein:

A is a direct bond, $-(CH_2)_a-$, $-(CH_2)_bCH=CH(CH_2)_c-$, or $-(CH_2)_bC\equiv C(CH_2)_c-$;

R_1 is aryl, heteroaryl or heterocycle fused to phenyl, each being optionally substituted with one to four substituents independently from R_3 ;

R_2 is $-R_3$, $-R_4$, $-(CH_2)_bC(=O)R_5$, $-(CH_2)_bC(=O)OR_5$,
 $-(CH_2)_bC(=O)NR_5R_6$, $-(CH_2)_bC(=O)NR_5(CH_2)_cC(=O)R_6$, $-(CH_2)_bNR_5C(=O)R_6$,
 $-(CH_2)_bNR_5C(=O)NR_6R_7$, $-(CH_2)_bNR_5R_6$, $-(CH_2)_bOR_5$, $-(CH_2)_bSO_dR_5$ or -
 $(CH_2)_bSO_2NR_5R_6$;

a is 1, 2, 3, 4, 5 or 6;

b and c are the same or different and at each occurrence independently 0, 1, 2, 3 or 4;

d is at each occurrence 0, 1 or 2;


R_3 is at each occurrence independently halogen, hydroxy, carboxy, alkyl, alkoxy, haloalkyl, acyloxy, thioalkyl, sulfinylalkyl, sulfonylalkyl, hydroxyalkyl, aryl, arylalkyl, heterocycle, heterocycloalkyl, $-C(=O)OR_8$, $-OC(=O)R_8$, $-C(=O)NR_8R_9$, $-C(=O)NR_8OR_9$, $-SO_2NR_8R_9$, $-NR_8SO_2R_9$, $-CN$, $-NO_2$, $-NR_8R_9$, $-NR_8C(=O)R_9$, $-NR_8C(=O)(CH_2)_bOR_9$, -
 $NR_8C(=O)(CH_2)_bR_9$, $-O(CH_2)_bNR_8R_9$, or heterocycle fused to phenyl;

R_4 is alkyl, aryl, arylalkyl, heterocycle or heterocycloalkyl, each being optionally substituted with one to four substituents independently from R_3 , or R_4 is halogen or hydroxy;

R_5 , R_6 and R_7 are the same or different and at each occurrence independently hydrogen, alkyl, aryl, arylalkyl, heterocycle or heterocycloalkyl, wherein each of R_5 , R_6 and R_7 are optionally substituted with one to four substituents independently selected from R_3 ; and


R_8 and R_9 are the same or different and at each occurrence independently hydrogen, alkyl, aryl, arylalkyl, heterocycle, or heterocycloalkyl, or R_8 and R_9 taken together with the atom or atoms to which they are bonded form a heterocycle, wherein each of R_8 , R_9 , and R_8 and R_9 taken together to form a heterocycle are optionally substituted with one to four substituents independently selected from R_3 .

11. (Withdrawn) The method of claim 2 wherein the compound has the following formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof.

12. (Withdrawn) The method of claim 3, wherein the compound has the following formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

wherein:

R₁ is aryl or heteroaryl optionally substituted with one to four substituents independently selected R₇;

R₂ is hydrogen;

R₃ is hydrogen or lower alkyl;

R₄ represents one to four optional substituents, wherein each substituent is the same or different and independently halogen, hydroxy, lower alkyl or lower alkoxy;

R₅ and R₆ are the same or different and independently -R₈, -(CH₂)_aC(=O)R₉, -(CH₂)_aC(=O)OR₉, -(CH₂)_aC(=O)NR₉R₁₀, -(CH₂)_aC(=O)NR₉(CH₂)_bC(=O)R₁₀, -(CH₂)_aNR₉C(=O)R₁₀, (CH₂)_aNR₁₁C(=O)NR₉R₁₀, -(CH₂)_aNR₉R₁₀, -(CH₂)_aOR₉, -(CH₂)_aSO_cR₉ or -(CH₂)_aSO₂NR₉R₁₀;

or R₅ and R₆ taken together with the nitrogen atom to which they are attached to form a heterocycle or substituted heterocycle;

R₇ is at each occurrence independently halogen, hydroxy, cyano, nitro, carboxy, alkyl, alkoxy, haloalkyl, acyloxy, thioalkyl, sulfinylalkyl, sulfonylalkyl, hydroxyalkyl, aryl, arylalkyl, heterocycle, heterocycloalkyl, -C(=O)OR₈, -OC(=O)R₈, -C(=O)NR₈R₉, -C(=O)NR₈OR₉, -SO_cR₈, -SO_cNR₈R₉, -NR₈SO_cR₉, -NR₈R₉, -NR₈C(=O)R₉, -NR₈C(=O)(CH₂)_bOR₉, -NR₈C(=O)(CH₂)_bR₉, -O(CH₂)_bNR₈R₉, or heterocycle fused to phenyl;


R₈, R₉, R₁₀ and R₁₁ are the same or different and at each occurrence independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, arylalkyl, heterocycle, heterocycloalkyl;

or R₈ and R₉ taken together with the atom or atoms to which they are attached to form a heterocycle;

a and *b* are the same or different and at each occurrence independently 0, 1, 2, 3 or 4; and

c is at each occurrence 0, 1 or 2.

13. (Withdrawn) The method of claim 3, wherein the compound has the following formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

wherein:

R₁ is aryl or heteroaryl optionally substituted with one to four substituents independently from R₇;

R₂ is hydrogen;

R₃ is hydrogen or lower alkyl;

R₄ represents one to four optional substituents, wherein each substituent is the same or different and independently halogen, hydroxy, lower alkyl or lower alkoxy;

R_5 and R_6 are the same or different and independently $-R_8$, $-(CH_2)_aC(=O)R_9$, $-(CH_2)_aC(=O)OR_9$, $-(CH_2)_aC(=O)NR_9R_{10}$, $-(CH_2)_aC(=O)NR_9(CH_2)_bC(=O)R_{10}$, $-(CH_2)_aNR_9C(=O)R_{10}$, $(CH_2)_aNR_{11}C(=O)NR_9R_{10}$, $-(CH_2)_aNR_9R_{10}$, $-(CH_2)_aOR_9$, $-(CH_2)_aSO_cR_9$ or $-(CH_2)_aSO_2NR_9R_{10}$;

or R_5 and R_6 taken together with the nitrogen atom to which they are attached to form a heterocycle or substituted heterocycle;

R_7 is at each occurrence independently halogen, hydroxy, cyano, nitro, carboxy, alkyl, alkoxy, haloalkyl, acyloxy, thioalkyl, sulfinylalkyl, sulfonylalkyl, hydroxyalkyl, aryl, arylalkyl, heterocycle, heterocycloalkyl, $-C(=O)OR_8$, $-OC(=O)R_8$, $-C(=O)NR_8R_9$, $-C(=O)NR_8OR_9$, $-SO_cR_8$, $-SO_cNR_8R_9$, $-NR_8SO_cR_9$, $-NR_8R_9$, $-NR_8C(=O)R_9$, $-NR_8C(=O)(CH_2)_bOR_9$, $-NR_8C(=O)(CH_2)_bR_9$, $-O(CH_2)_bNR_8R_9$, or heterocycle fused to phenyl;


R_8 , R_9 , R_{10} and R_{11} are the same or different and at each occurrence independently hydrogen, alkyl, aryl, arylalkyl, heterocycle, heterocycloalkyl;

or R_8 and R_9 taken together with the atom or atoms to which they are attached to form a heterocycle;

a and b are the same or different and at each occurrence independently 0, 1, 2, 3 or 4; and

c is at each occurrence 0, 1 or 2.

14. (Withdrawn) The method of claim 3, wherein the compound has the following formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

wherein:

R₁ is aryl or heteroaryl optionally substituted with one to four substituents independently from R₇;

R₂ is hydrogen;

R₃ is hydrogen or lower alkyl;

R₄ represents one to four optional substituents, wherein each substituent is the same or different and independently from halogen, hydroxy, lower alkyl or lower alkoxy;

R₅ and R₆ are the same or different and independently -R₈, -(CH₂)_aC(=O)R₉, -(CH₂)_aC(=O)OR₉, -(CH₂)_aC(=O)NR₉R₁₀, -(CH₂)_aC(=O)NR₉(CH₂)_bC(=O)R₁₀, -(CH₂)_aNR₉C(=O)R₁₀, (CH₂)_aNR₁₁C(=O)NR₉R₁₀, -(CH₂)_aNR₉R₁₀, -(CH₂)_aOR₉, -(CH₂)_aSO_cR₉ or -(CH₂)_aSO₂NR₉R₁₀;

or R₅ and R₆ taken together with the nitrogen atom to which they are attached to form a heterocycle;

R₇ is at each occurrence independently halogen, hydroxy, cyano, nitro, carboxy, alkyl, alkoxy, haloalkyl, acyloxy, thioalkyl, sulfinylalkyl, sulfonylalkyl, hydroxyalkyl, aryl, arylalkyl, heterocycle, heterocycloalkyl, -C(=O)OR₈, -OC(=O)R₈, -C(=O)NR₈R₉, -C(=O)NR₈OR₉, -SO_cR₈, -SO_cNR₈R₉, -NR₈SO_cR₉, -NR₈R₉, -NR₈C(=O)R₉, -NR₈C(=O)(CH₂)_bOR₉, -NR₈C(=O)(CH₂)_bR₉, -O(CH₂)_bNR₈R₉, or heterocycle fused to phenyl;

R₈, R₉, R₁₀ and R₁₁ are the same or different and at each occurrence independently hydrogen, alkyl, substituted alkyl, aryl, arylalkyl, heterocycle, heterocycloalkyl;

or R₈ and R₉ taken together with the atom or atoms to which they are attached to form a heterocycle;


a and *b* are the same or different and at each occurrence independently 0, 1, 2, 3 or 4; and

c is at each occurrence 0, 1 or 2.

15. (Withdrawn) The method of claim 4, wherein R₀ is -O-.

16. (Withdrawn) The method of claim 4, wherein R_0 is -S-.
17. (Withdrawn) The method of claim 4, wherein R_0 is -S(O)-.
18. (Withdrawn) The method of claim 4, wherein R_0 is -S(O)₂-.
19. (Withdrawn) The method of claim 4, wherein R_0 is NH.
20. (Withdrawn) The method of claim 04, wherein R_0 is CH₂-.

21. (Withdrawn) The method of claim 4, wherein the compound has the following formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof.

22. (Original) The method of claim 1, further comprising administering a second active agent.
23. (Original) The method of claim 2, further comprising administering a second active agent.
24. (Withdrawn) The method of claim 3, further comprising administering a second active agent.
25. (Withdrawn) The method of claim 4, further comprising administering a second active agent.
26. (Original) The method of claim 22, wherein the second active agent is an antidepressant, antihypertensive, anxiolytic, calcium channel blocker, muscle relaxant, non-narcotic analgesic, anti-inflammatory agent, cox-2 inhibitor, alpha-adrenergic receptor

agonist or antagonist, ketamine, anesthetics, immunomodulatory agent, immunosuppressive agent, corticosteroid, hyperbaric oxygen, anticonvulsant, an IMiD[®], a SelCID[®], or a combination thereof.

27. (Original) The method of claim 22, wherein the second active agent is gabapentin, thalidomide, salicylic acid acetate, ketamine, celocoxib, carbamazepine, oxcarbazepine, phenytoin, sodium valproate, prednisone, nifedipine, clonidine, oxycodone, meperidine, morphine sulfate, hydromorphone, fentanyl, acetaminophen, ibuprofen, naproxen sodium, griseofulvin, amitriptyline, imipramine, doxepin, or a pharmaceutically acceptable salt, solvate or stereoisomer thereof.

28. (Original) The method of claim 1, wherein the pain is complex regional pain syndrome.

29. (Original) The method of claim 28, wherein the complex regional pain syndrome is type I or type II.

30. (Previously Presented) The method of claim 28, wherein the complex regional pain syndrome is stage I, stage II or stage III of complex regional pain syndrome type I.

31. (Currently Amended) The method of claim 1, wherein the pain is or is associated with autonomic dysfunction, trigeminal neuralgia, post-herpetic neuralgia, cancer-related pain, phantom limb pain, fibromyalgia, chronic fatigue syndrome, radiculopathy, inability to initiate movement, weakness, tremor, muscle spasm, dystonia, dystrophy, atrophy, edema, stiffness, joint tenderness, increased sweating, sensitivity to temperature, light touch (allodynia), color change to the skin, hyperthermic or hypothermic, increased nail and hair growth, early bony changes, hyperhidrotic with livedo reticularis or cyanosis, lost hair, ridged, cracked or brittle nails, dry hand, diffuse osteoporosis, irreversible tissue damage, thin and shiny skin, joint contractures, marked bone demineralization, diabetic neuropathy, luetic neuropathy, or painful neuropathy induced iatrogenically by a drug, ~~or another painful neuropathic condition.~~

32. (Original) The method of claim 1, wherein the pain is nociceptive pain.

33. (Original) The method of claim 32, wherein the nociceptive pain is associated with a cut or contusion of the skin; a chemical or thermal burn; osteoarthritis; rheumatoid arthritis; or tendonitis.
34. (Original) The method of claim 1, wherein the pain is neuropathic pain.
35. (Original) The method of claim 34, wherein the neuropathic pain is associated with stroke, diabetic neuropathy, luetic neuropathy, postherpetic neuralgia, trigeminal neuralgia, fibromyalgia, or painful neuropathy induced iatrogenically by a drug.
36. (Withdrawn) A pharmaceutical composition comprising an effective amount of a JNK Inhibitor and an antidepressant, antihypertensive agent, anxiolytic agent, calcium channel blocker, muscle relaxant, non-narcotic analgesic, anti-inflammatory agent, cox-2 inhibitor, alpha-adrenergic receptor agonist or antagonist, ketamine, an anesthetic, an immunomodulatory agent, an immunosuppressive agent, a corticosteroid, hyperbaric oxygen, an anticonvulsant, an IMiD[®], a SelCID[®], or a combination thereof.